

15th Water-Rock Interaction International Symposium, WRI-15

Environmental tracers in groundwaters and porewaters to understand groundwater movement through an argillaceous aquitard

Stacey C. Priestley^{a,1}, Andrew J. Love^a, Vincent Post^b, Paul Shand^c, Daniel Wohling^d,
Rolf Kipfer^e, Timothy E. Payne^f, Martin Stute^g, Lina Tyroller^e

^aFlinders University and NCGRT, Bedford Park, SA 5042, Australia

^bFederal Institute for Geosciences and Natural (BGR), Geozentrum Hannover, Stilleweg 2, D-30655, Germany

^cCSIRO land and water, Waite Campus, Glen Osmond SA 5064, Australia

^dDEWNR, L11 25 Grenfell Street, Adelaide, SA 5000, Australia

^eSwiss Federal Institute of Aquatic Science and Technology, 8600 Duebendorf, Switzerland and Swiss Federal Institute of Technology, 8092 Zurich, Switzerland

^fANSTO, Lucas Heights Campus, NSW 2234, Australia

^gLamont-Doherty Earth Observatory, 61 route 9W, Palisades, NY 10964, USA

Abstract

Inter-aquifer leakage through an aquitard can be an important component of groundwater flow and can occur by diffuse leakage or via preferential pathways along secondary permeability features. In order to properly characterize inter-aquifer leakage it is desirable to use both porewater from the aquitard in conjunction with a regional investigation of groundwater in the aquifers. The aim of this study was to characterize inter-aquifer leakage through a regionally extensive aquitard between the Great Artesian Basin (GAB) and the deeper Arckaringa Basin of Australia. Chloride concentrations in the aquitard porewater profile indicates that transport through the aquitard is dominated by diffusion, but there is evidence at t least one location for increased interaquifer leakage due to secondary permeability features.

© 2017 Published by Elsevier B.V. This is an open access article under the CC BY-NC-ND license

(<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Peer-review under responsibility of the organizing committee of WRI-15

Keywords: inter-aquifer leakage; aquitard; environmental tracers

* Corresponding author. Tel.: +61 8 8201 5302.

E-mail address: stacey.priestley@flinders.edu.au

1. Introduction

Aquitards are very important, yet poorly understood components in groundwater flow systems¹. Regionally extensive aquitards confine and separate aquifers and play an important role in the physical and chemical evolution of groundwater, and can provide water from storage to pumped aquifers^{1,2}. Groundwater movement through an aquitard, here referred to as inter-aquifer leakage, can be an important component of groundwater flow, although it is orders of magnitude slower than through aquifers³. Quantification of inter-aquifer leakage is important for groundwater resource evaluation and extraction industries including unconventional gas and ore extraction.

Inter-aquifer leakage through a laterally-extensive aquitard can be severely limited but even in the tightest clay or shale aquitard there is always a small amount of diffuse leakage². The inter-aquifer leakage rate through the aquitard can increase if there are secondary permeability features through the aquitard^{2,4-6}.

To quantify inter-aquifer leakage different techniques are available. Chemical profiles through the aquitard can be used to investigate the transport mechanisms in the aquitard pore water^{5,7-10}, whereas detection of inter-aquifer leakage through secondary permeability features requires a more regional scale study. Inter-aquifer leakage through secondary permeability features has been identified in some regional groundwater investigations using hydraulic head measurements, environmental tracers and dating tracers in aquifers¹¹⁻¹⁵.

Several authors³ urged the use of and comparison of methods at different scales within the same formation. The aim of this study was to investigate inter-aquifer leakage through a regionally extensive aquitard between the Great Artesian Basin (GAB) and the Arckaringa Basin.

2. Study area

This study investigated inter-aquifer leakage between the GAB and Arckaringa Basin in the central to far north region of South Australia (Fig. 1, inset). The GAB comprises Jurassic to Cretaceous sediments that overlie the Arckaringa Basin which is a Late Carboniferous to Early Permian sedimentary basin. The main aquifer unit in the GAB is the J-K aquifer and in the study area the main aquifer in the Arckaringa Basin is the Boorthanna Formation (Figure 1a). The J-K aquifer and Boorthanna Formation are separated by a mudstone, siltstone and shale aquitard, known as the Stuart Range Formation (Figure 1b).

3. Methods

To investigate inter-aquifer leakage in between the GAB and Arckaringa Basin regional groundwater samples as well as samples from the aquitard were collected. A core through the aquitard was drilled and aquitard samples collected for environmental tracer analysis¹⁰. Regional groundwater samples were analysed for various environmental tracers and isotopes including major elements, stable isotopes of water, ¹⁴C, ³⁶Cl, ⁸⁷Sr/⁸⁶Sr, uranium isotopes, as well as noble gases.

4. Results and discussion

The chloride concentrations in the aquitard porewater profile are shown in Figure 2. Modelling the profile using the one-dimensional advective-dispersive equation showed that the dominant transport through the aquitard is diffusion. The regional groundwater samples show evidence of inter-aquifer leakage through secondary permeability features in the Stuart Range Formation in the center of the basin. This inter-aquifer leakage has been identified by overlap in Sr and stable water isotope values, chloride concentrations as well as other hydrochemical evidence of mixing with shallower groundwater with shorter residence times. Further evidence is provided by head data which showed a slight draw-down in the upper aquifer during pumping in the lower aquifer.

The results show that flow through the aquitard is dominated by diffusion; there is at least one location with increased inter-aquifer leakage due to secondary permeability features. These results are similar to other studies which have found inter-aquifer leakage through preferential pathways through aquitards^{11,15}.

5. Conclusions

These results show that diffusion, although dominant is not the only flow mechanism through regional scale aquitards, but secondary permeability features such as fractures and discontinuities can contribute to inter-aquifer leakage. Although cross formational flow is not regionally extensive, it may have a disproportionate impact on water flow and water quality.




Fig. 1. (a) Map showing upper aquifer extent in green and the density corrected potentiometric surface in green dashed lines. The lower aquifer extent is shown in blue with the density corrected potentiometric surface in blue dashed lines. Cross-section through the study area is indicated with a solid black line (b) stratigraphy of hydrogeological cross-section.




Fig. 2. Chloride concentration through the aquitard.

References

1. Back W. Role of aquitards in hydrogeochemical systems: a synopsis. *Appl Geochem*, 1986;1:427-437.
2. Cherry JA, Parker BL., *Role of Aquitards in the Protection of Aquifers from Contamination: A "State of Science" Report*. Denver, USA: AWWA Research Foundation Vol. 47; 2004. p. 15-21.
3. Battle-Aguilar J, Cook PG, Harrington GA. Comparison of hydraulic and chemical methods for determining hydraulic conductivity and leakage rates in argillaceous aquitards. *Journal of Hydrology*, 2016;532:102-121.
4. Neuzil CE. Groundwater-flow in low-permeability environments. *Water Resources Research*, 1986;22:1163-1195.
5. Hendry MJ, Wassenar LI. Transport and geochemical controls on the distribution of solutes and stable isotopes in a thick clay-rich till aquitard, Canada. *Isotopes in Environmental and Health Studies*, 2004;40:3-19.
6. Hart DJ, Bradbury KR, Feinstein DT. The vertical hydraulic conductivity of an aquitard at two spatial scales. *Ground Water*, 2006;44:201-211.
7. Desaulniers DE, Cherry JA. Origin and movement of groundwater and major ions in a thick deposit of Champlain sea clay near Montreal. *Canadian Geotechnical Journal*, 1989;26:80-89.
8. Remenda VH, van der Kamp G, Cherry JA. Use of vertical profiles of delta O-18 to constrain estimates of hydraulic conductivity in a thick, unfractured aquitard. *Water Resources Research*, 1996;32:2979-2987.
9. Gimmi T, Waber HN, Gautschi A, Rubel A. Stable water isotopes in pore water of Jurassic argillaceous rocks as tracers for solute transport over large spatial and temporal scales. *Water Resources Research*, 2007;43:1-16.
10. Mazurek M, Alt-Epping P, Bath A, Gimmi T, Waber HN. *Natural Tracer Profiles Across Argillaceous Formations: The CLAYTRAC Project*. OECD Publishing; 2009. p. 136-150.
11. Love AJ, Herczeg AL, Walker G. Transport of water and solutes across a regional aquitard inferred from deuterium and chloride profiles Otway Basin, Australia. *Isotopes in Water Resources Management (Proc Symp Vienna, 1995)*: IAEA; 1996. p. 273-286.
12. Clark JF, Stute M, Schlosser P, Drenkard S. A tracer study of the Floridan aquifer in southeastern Georgia: Implications for groundwater flow and paleoclimate. *Water Resour Res*. 1997;33:281-289.
13. Dogramaci SS, Herczeg AL. Strontium and carbon isotope constraints on carbonate-solution interactions and inter-aquifer mixing in groundwaters of the semi-arid Murray Basin, Australia. *J Hydrol*. 2002;262:50-67.
14. Althaus R, Klump S, Onnis A, Kipfer R, Purtschert R, Stauffer F, Kinzelbach W. Noble gas tracers for characterisation of flow dynamics and origin of groundwater: A case study in Switzerland. *J Hydrol*. 2009;370:64-72.
15. Gardner PM, Heilwell VM. A multiple-tracer approach to understanding regional groundwater flow in the Snake Valley area of the Eastern Great Basin, USA. *Appl Geochem*. 2014;45:33-49.